

Research paper

# Differentiation of buckwheat cultivars (*Fagopyrum esculentum* Moench) by yield structure elements and productivity under the conditions of the Forest-Steppe of Ukraine

Dmytro CHUIKO<sup>1\*</sup>, Vladyslav STRYZHAK<sup>1</sup>, Oleh TRYHUB<sup>2</sup>

<sup>1</sup> State Biotechnological University, Department of Genetics, Breeding and Seed Production 61002, 44 Alchevskykh Str., Kharkiv, Ukraine.

<sup>2</sup> Ustymivska Experimental Station of Plant Production of Plant Production Institute nd. V.Ya. Yuryev of NAAS of Ukraine, Ustymivka, Poltavska reg., Ukraine.

\* Corresponding author

**E-mail addresses of authors:** chuiko93ua@gmail.com (DC); luch6625@gmail.com (VS); trygub\_oleg@ukr.net (OT)

DOI: <https://doi.org/10.3986/fag0054>

**Key words:** buckwheat; cultivar; breeding; productivity; thousand-kernel weight; cluster analysis.

**Received:** February 1, 2026; accepted May 30, 2026

## ABSTRACT

The results of a three-year research (2023–2025) of 31 common buckwheat varieties (*Fagopyrum esculentum* Moench) of different ecological and geographical origins under the conditions of the Eastern Forest-Steppe of Ukraine are presented. The peculiarities of the formation of plant height, number of leaves and inflorescences, plant productivity, thousand-kernel weight, number of kernels per plant, and kernel set were investigated. The determining role of growth type in shaping plant architecture and productivity was established.

It was found that productivity is formed mainly due to the number of kernels per plant and kernel set ( $r = 0.95$ ), whereas thousand-kernel weight is a relatively stable trait ( $V = 7.5\%$ ). According to the results of k-means clustering, a group of intensive-type varieties was identified, characterized by the highest average values within the cluster for plant productivity (2.4 g), number of kernels (85.7 pcs.), thousand-kernel weight (27.9 g), and kernel set (3.8).

The varieties 'Pidmoskovna T', 'M NF-84B', 'Mariia', 'Sumchanka', 'Roksolana', 'Krupynka', 'Selianochka', and 'Simka' were identified as promising sources of valuable traits for further breeding aimed at improving productivity and thousand-kernel weight.

## INTRODUCTION

Effective common buckwheat (*Fagopyrum esculentum* Moench) breeding requires the availability of broad genetic diversity preserved in buckwheat germplasm collections worldwide. Global buckwheat germplasm has demonstrated considerable variation in morphological and agronomic traits (Janovská et al., 2021; Rauf et al., 2020; Ghiselli et al., 2016). This diversity represents a fundamental resource for breeding programs, as genetic variability is the primary driver of selection and evolution and serves as the genetic basis for breeders (Dutta et al., 2025; Bhandari et al., 2017).

A detailed study of the morphological and genetic diversity of germplasm is indispensable for identifying genomic regions controlling important agronomic traits and for detecting valuable genetic accessions carrying favorable alleles that can be used in marker-assisted breeding programs (Pipan et al., 2023). Buckwheat varieties that have undergone production testing and have been cultivated in different countries are of particular value as breeding material, since their performance can serve as a reliable basis for comparative evaluation.

Significant progress in global buckwheat breeding practice has been achieved in addressing key agronomic problems through conventional breeding methods. Improvement of individual traits occurs gradually, and in Japan, varieties resistant to lodging such as ‘Tachikane’ and ‘Nijiyutaka’, as well as a variety resistant to pre-harvest sprouting, ‘Harunoyuki’, have been developed (Ohsawa, 2020). Modern breeding efforts have also demonstrated success in developing regionally adapted, high-yielding varieties for Ukraine’s conditions, including ‘Oranta’, ‘Ruslan’, and ‘Volia’ (Shakalii et al., 2023). These achievements of classical breeding illustrate effective solutions to fundamental production constraints that previously limited buckwheat acreage and yield.

Despite achievements in buckwheat breeding, several significant challenges remain in developing modern varieties. The main issues include insufficient understanding of the inheritance patterns of most economically valuable traits related to yield and the synthesis of biologically active compounds, low seed productivity, seed shattering, uneven flowering, and limited knowledge of gene functioning. Prospects for further breeding include reducing polyphenol content through selection and developing varieties with improved processing and storage characteristics. In general, breeding efforts aim to improve buckwheat’s agronomic traits through convention-

al breeding combined with biotechnological methods (Luthar et al., 2021; Chuiko & Tryhub, 2025; Suzuki et al., 2020).

Thus, the main objective of our study was to evaluate a collection of diploid common buckwheat varieties (*F. esculentum* Moench) of different ecological and geographical origins obtained from the Plant Genetic Resources Bank of Ukraine for the expression of key agronomically valuable traits under the conditions of the Eastern Forest-Steppe of Ukraine.

## MATERIAL AND METHODS

Field and laboratory studies of a collection of diploid common buckwheat varieties (*F. esculentum* Moench) were conducted during 2023–2025 at the experimental field of the Department of Genetics, Breeding and Seed Production of the State Biotechnological University. The experimental fields are located in the central part of Kharkiv region (Dokuchaievske settlement, Ukraine) at the boundary of the Forest-Steppe and Steppe natural-climatic zones (geographical coordinates: 49.902144, 36.446238).

The experimental plots were established according to standard breeding methodologies within a breeding crop rotation. The predecessor was bare fallow. Sowing was



**Figure 1.** Buckwheat variety *Bashkirska Krasnostebelny* (UC0102179).

**Table 1.** Characteristics of the studied common buckwheat varieties (*F. esculentum* Moench.).

| No | National Catalogue ID | Region and country of origin   | Variety name             | Growth type* |
|----|-----------------------|--------------------------------|--------------------------|--------------|
| 1  | UC0100246             | Primorsky Kray (RU)            | Prymorska 5              | Indet.       |
| 2  | UC0100261             | Primorsky Kray (RU)            | Mariia                   | Indet.       |
| 3  | UC0100989             | Minsk Region (BY)              | Biloruska Skorostyhl'a   | Indet.       |
| 4  | UC0101130             | Oryol Oblast (RU)              | Kozachka                 | Indet.       |
| 5  | UC0100991             | Minsk Region (BY)              | Biloruska Odnostebelna   | Indet.       |
| 6  | UC0101006             | Oryol Oblast (RU)              | Krupynka                 | Det.         |
| 7  | UC0101083             | Poltavska Oblast (UA)          | Pidmoskovna T            | Indet.       |
| 8  | UC0100352             | Khmelnyskyi Oblast (UA)        | M NF-84B                 | Indet.       |
| 9  | UC0100362             | Minsk Region (BY)              | Biloruska Homostyl'na    | Indet.       |
| 10 | UC0100365             | Minsk Region (BY)              | Chornohlazka             | Indet.       |
| 11 | UC0100367             | Minsk Region (BY)              | Zhniarka                 | Indet.       |
| 12 | UC0100501             | Sumska Oblast (UA)             | Yampolska Mistseva       | Indet.       |
| 13 | UC0101981             | Sumska Oblast (UA)             | Yuvileina 100            | Det.         |
| 14 | UC0101987             | Minsk Region (BY)              | Karmen                   | Det.         |
| 15 | UC0100942             | Khmelnyskyi Oblast (UA)        | Exida                    | Indet.       |
| 16 | UC0101189             | Minsk Region (BY)              | Anita Biloruska          | Indet.       |
| 17 | UC0101200             | Khmelnyskyi Oblast (UA)        | Viktoriia Podilska       | Indet.       |
| 18 | UC0101198             | Khmelnyskyi Oblast (UA)        | Roksolana                | Indet.       |
| 19 | UC0101190             | Minsk Region (BY)              | Zhaleika                 | Indet.       |
| 20 | UC0101197             | Minsk Region (BY)              | Smuhlianka               | Det.         |
| 21 | UC0101977             | Japan (JP)                     | Ketavase                 | Indet.       |
| 22 | UC0102165             | Minsk Region (BY)              | Kvietka Determinantna    | Det.         |
| 23 | UC0102166             | Minsk Region (BY)              | Charovnit'sa             | Indet.       |
| 24 | UC0102186             | Sumska Oblast (UA)             | Sumchanka                | Det.         |
| 25 | UC0102179             | Republic of Bashkortostan (RU) | Bashkyska Krasnostebelna | Indet.       |
| 26 | UC0102193             | Minsk Region (BY)              | Vlada                    | Det.         |
| 27 | UC0102203             | Minsk Region (BY)              | Amethyst                 | Indet.       |
| 28 | UC0102204             | Minsk Region (BY)              | Lakneia                  | Det.         |
| 29 | UC0100206             | Sumska Oblast (UA)             | Selianochka              | Indet.       |
| 30 | UKR008:01705          | Sumska Oblast (UA)             | Simka                    | Indet.       |
| 31 | UC0102228             | Sumska Oblast (UA)             | Oksana                   | Indet.       |

\* – Growth type: Indeterminate type – Indet.; Determinate type – Det.

carried out in the second decade of May. Each variety was grown in four replications. The seeding rate was 80 viable seeds per linear meter with row spacing of 45 cm. The arrangement of replications was systematic (Ermantraut et al., 2014). No additional fertilization or herbicide application was performed. According to the FAO soil classification, the soil of the experimental field is defined as Calcic Voronic Chernozem CL UE1 (Jahn et al., 2006).

The research material consisted of 31 varieties obtained from the Ustymivka Experimental Station of Plant Production of the V.Ya. Yuryev Institute of Plant Produc-

tion of NAAS of Ukraine. The collection includes buckwheat varieties of different ecological and geographical origins. A detailed description of the studied common buckwheat accessions is presented in Table 1 (Fig. 1).

Observations and measurements in the experimental plots were carried out in accordance with the "Methodological guidelines for studying collection accessions of maize, sorghum, and cereal crops (millet, buckwheat, rice)" (Shmaryayev et al., 1968) and the "Methodology of state variety testing of agricultural crops (cereals, pseudocereals, and legumes)" (Volkodav, 2001).

Morphological and productivity traits were evaluated on 20 representative plants in each replication. Plant height was measured from the soil surface to the plant apex. The number of leaves and inflorescences per plant was determined by direct counting. Plant productivity was assessed through manual threshing followed by weighing, and the mean value was calculated for each replication. The seed set index was determined as the ratio of the average number of seeds per plant to the average number of inflorescences per plant within a replication.

In 2023, temperature conditions during the buckwheat growing season were close to the long-term average. In May and June, a slight decrease in temperature was observed (−0.2 and −0.4 °C, respectively), whereas in July, August, and September, temperatures exceeded the norm by 0.8, 2.7, and 2.3 °C, respectively.

In 2024, warmer conditions prevailed throughout the growing season. May was slightly cooler (−0.8 °C), but from June to September, a significant excess over the long-term averages was recorded: +2.6 °C in June, +4.5 °C in July, +2.9 °C in August, and +5.3 °C in September.

The temperature regime in 2025 was moderate, close to the long-term average. In May and June, temperatures were below the norm by 0.9 and 0.6 °C, in July they exceeded it by 3.0 °C, in August they were close to the norm (−0.3 °C), and in September they were higher by 1.3 °C.

In 2023, the distribution of precipitation during the buckwheat growing season was contrasting: moisture deficit in May and June (−13.0 and −17.3 mm from the norm) was followed by excessive rainfall in July, when 153.8 mm of precipitation fell, which was 2.5 times higher than the long-term average. In August and September, a precipitation deficit was again observed (−13.2 and −18.5 mm).

In 2024, a pronounced drought occurred: only 92.1 mm of precipitation fell during the growing season, which is about one-third of the norm. All months were characterized by a significant moisture deficit, with July being the most critical (−47.0 mm), and no precipitation was recorded in September.

In 2025, moisture conditions were better than in 2024 and generally close to the norm. In May, July, and August, precipitation exceeded the long-term averages by 21.8, 12.8, and 9.9 mm, respectively, while in June and September a deficit was observed (−29.4 and −21.8 mm). Overall, during the growing season, total precipitation was 6.7 mm below the long-term average.

Statistical data processing was performed using PAST 5.3 software (Hammer & Harper, 2001). To characterize the studied varieties, cluster analysis with Euclidean distances (k-means clustering) was applied. The relationships between traits were evaluated using Pearson's linear correlation.

To normalize the values for cluster analysis, the Z-score standardization method (standardization by mean and standard deviation) was applied according to the formula (Cheadle et al., 2003):

$$z = \frac{x - \mu}{\sigma}$$

where:

z — standardized (normalized) value;

x — original value;

μ — arithmetic mean of the trait;

σ — standard deviation of the trait.


## RESULTS AND DISCUSSION

Plant height is an important technological and breeding trait in buckwheat. This trait can vary considerably across plant genotypes. The main factors influencing plant height are growth type (determinate or indeterminate) and the length of the growing period, which may be associated with the genotype's ecological and geographical origin.

According to our field studies conducted in 2023–2025, growth type (determinate or indeterminate) was the primary factor determining plant height in the studied buckwheat varieties. The greatest plant height was observed in indeterminate varieties such as 'Ketavase' – 114.3 ± 14.0 cm, 'M NF-84B' – 104.9 ± 11.1 cm, 'Mariia' – 102.6 ± 18.2 cm, 'Viktoriia Podilska' – 102.1 ± 8.7 cm, and 'Roksolana' – 102.1 ± 2.2 cm.

Conversely, the lowest plant height over the years of study was recorded in determinate varieties, for example 'Vlada' – 74.7 ± 3.8 cm, 'Lakneia' – 78.7 ± 9.6 cm, and 'Krupynka' – 83.3 ± 13.4 cm (Figure 2). Similar patterns in plant height distribution have been reported by other authors under different climatic conditions (Amelin et al., 2020; Kharchenko & Tryhub, 2018; Kasajima et al., 2016).

Buckwheat productivity is influenced by numerous interrelated plant characteristics that affect both photosynthetic capacity and seed yield. The leaf system serves as the main photosynthetic apparatus, converting solar energy into resources that support growth and seed for-



**Figure 2.** Formation of height traits in buckwheat varieties, average for the years of study.

mation. Buckwheat exhibits an indeterminate growth pattern, continuously forming both leaves and flowers during most of the growing season, which makes the balance between vegetative and reproductive structures particularly important for optimizing overall plant productivity. In general, the ratio of the number of leaves to the number of inflorescences in buckwheat (*F. esculentum* Moench) is approximately 1:1 (Alekseyeva et al., 2004). The main problem is the insufficient supply of photosynthetic surface to support a large number of inflorescences, which leads to excessive abortion of seed ovaries at early stages of development.


According to our results, the number of leaves and the number of inflorescences per plant varied considerably depending on the year of cultivation. The highest average values over the years were observed in the varieties 'Prymorska 5', 'Mariia', 'Biloruska Skorostyhl'a', 'Kozachka', and 'Biloruska Homostylna', ranging from 41.1 to 48.5 leaves and from 30.1 to 35.9 inflorescences per plant, respectively.

At the same time, low-growing varieties such as 'Bashkyrska Krasnostebelna', 'Vlada', 'Amethyst', 'Lakneia', and 'Selianochka' proved to be the most efficient

in terms of plant architecture and leaf support of inflorescences, with average values of 24.8–31.2 leaves and 14.6–20.6 inflorescences per plant, respectively. Thus, these varieties exhibited the highest leaf-to-inflorescence ratio, ranging from 1.5 to 1.7 leaves per inflorescence. In other studies, a reduction in the number of inflorescences combined with an increase in leaf number has been shown to enhance plant productivity and overall buckwheat yield (Hornýák et al., 2020). In the other studied varieties, the leaf supply per inflorescence ranged from 1.2 to 1.5 (Fig. 3).

The generalized results of the field experiments indicate substantial genotypic variability among the studied buckwheat cultivars. The mean productivity per plant was 1.7 g, ranging from 1.1 to 3.2 g; thousand-kernel weight averaged 27.0 g (23.1–31.5 g); the number of kernels per plant was 63.5 (43.4–112.7); and the seed set index was 3.0 (1.4–4.7). The coefficient of variation was highest for productivity per plant (26.2%), number of kernels per plant (24.1%), and seed set index (27.1%), indicating a high sensitivity of these traits to genotype and growing conditions, whereas thousand-kernel weight was relatively stable ( $V = 7.5\%$ ).





**Figure 3.** Characteristics of leaf and inflorescence formation in buckwheat varieties, multi-year average.

The highest productivity per plant was recorded for the cultivar 'Pidmoskovna T' ( $3.2 \pm 1.9$  g), which was accompanied by the maximum number of kernels per plant ( $112.7 \pm 56.2$ ) and the highest seed set index ( $4.7 \pm 2.1$ ). High values of productivity and kernel number were also observed in the cultivars 'M NF-84B' ( $2.6 \pm 0.6$  g;  $82.7 \pm 12.3$  kernels), 'Mariia' ( $2.3 \pm 0.7$  g;  $80.5 \pm 21.4$  kernels), and 'Sumchanka' ( $2.3 \pm 1.1$  g;  $83.2 \pm 39.7$  kernels). These cultivars combine increased seed productivity with a relatively large number of formed kernels, indicating efficient realization of the generative potential of the plants.

Cultivars with the lowest productivity per plant — 'Exida' ( $1.1 \pm 0.4$  g), 'Ketavase' ( $1.2 \pm 0.4$  g), 'Biloruska Skorostyhl'a' ( $1.3 \pm 0.3$  g), and 'Biloruska Odnostebelna' ( $1.3 \pm 0.5$  g) — were also characterized by a comparatively lower number of kernels per plant (43.4–49.4), demonstrating a direct dependence of productivity on the quantitative parameters of generative organs and a very strong correlation between productivity per plant and kernel number ( $r = 0.95$ ).

Thousand-kernel weight varied within a relatively narrow range; however, cultivars with larger kernels were clearly distinguished, including 'M NF-84B' ( $31.5 \pm 2.5$

g), 'Roksolana' ( $30.7 \pm 1.8$  g), 'Krupynka' and 'Selianochka' (29.6 g each), 'Simka' ( $29.3 \pm 1.1$  g), and 'Kozachka' ( $29.0 \pm 2.8$  g). In contrast, the lowest thousand-kernel weight was observed in 'Exida' ( $23.1 \pm 0.5$  g), 'Yampolska Mistseva' ( $24.2 \pm 0.9$  g), 'Bashkyrska Krasnostebelna', and 'Charovnit'sa' ( $24.5 \pm 1.0$  g), which may limit their value for groat production but does not necessarily reduce overall productivity.

The seed set index, which reflects the ratio between the number of kernels and inflorescences, showed pronounced differentiation among the cultivars. The highest values were recorded for 'Pidmoskovna T' ( $4.7 \pm 2.1$ ), 'Vlada' ( $4.6 \pm 2.1$ ), 'Zhniarka' ( $4.1 \pm 1.7$ ), 'Lakneia' ( $3.9 \pm 1.7$ ), 'Bashkyrska Krasnostebelna' ( $3.9 \pm 1.7$ ), and 'Smuglianka' ( $3.6 \pm 1.4$ ). This indicates a more efficient realization of fertilization potential and a lower level of ovary abortion. In contrast, the lowest seed set indices were observed in 'Biloruska Skorostyhl'a' ( $1.4 \pm 0.3$ ), 'Ketavase' ( $1.5 \pm 0.6$ ), and 'Kozachka' ( $1.5 \pm 0.7$ ), suggesting less efficient seed formation under a comparable number of inflorescences per plant.

Overall, the analysis of yield structure demonstrates that high productivity in buckwheat is primarily deter-


mined by an increase in the number of kernels per plant and a high level of seed set, whereas thousand-kernel weight plays a stabilizing but secondary role. The identified cultivars 'Pidmoskovna T', 'M NF-84B', 'Mariia', and

'Sumchanka' represent valuable sources of seed productivity traits, while 'M NF-84B', 'Roksolana', 'Krupynka', and 'Simka' are important sources of large seed size for further buckwheat breeding programs (Table 2).

**Table 2.** Characteristics of buckwheat varieties by yield component traits, multi-year average.

| Cultivar                  | Productivity per plant, g., $\pm$ SD* | Thousand-kernel weight, g., $\pm$ SD | Number of kernels per plant, pcs., $\pm$ SD | Number of kernels per inflorescence, $\pm$ SD |
|---------------------------|---------------------------------------|--------------------------------------|---------------------------------------------|-----------------------------------------------|
| Prymorska 5               | 1.7 $\pm$ 0.4                         | 26.7 $\pm$ 2.8                       | 63.5 $\pm$ 14.0                             | 2.1 $\pm$ 0.1                                 |
| Mariia                    | 2.3 $\pm$ 0.7                         | 28.7 $\pm$ 0.8                       | 80.5 $\pm$ 21.4                             | 2.6 $\pm$ 1.3                                 |
| Biloruska Skorostyhla     | 1.3 $\pm$ 0.3                         | 27.2 $\pm$ 2.3                       | 46.8 $\pm$ 7.1                              | 1.4 $\pm$ 0.3                                 |
| Kozachka                  | 1.4 $\pm$ 0.7                         | 29.0 $\pm$ 2.8                       | 47.3 $\pm$ 17.8                             | 1.5 $\pm$ 0.7                                 |
| Biloruska Odnostebelna    | 1.3 $\pm$ 0.5                         | 26.5 $\pm$ 1.3                       | 49.4 $\pm$ 17.7                             | 3.0 $\pm$ 1.2                                 |
| Krupynka                  | 1.8 $\pm$ 0.7                         | 29.6 $\pm$ 3.6                       | 60.9 $\pm$ 19.2                             | 3.0 $\pm$ 1.3                                 |
| Pidmoskovna T             | 3.2 $\pm$ 1.9                         | 27.5 $\pm$ 2.6                       | 112.7 $\pm$ 56.2                            | 4.7 $\pm$ 2.1                                 |
| M NF-84B                  | 2.6 $\pm$ 0.6                         | 31.5 $\pm$ 2.5                       | 82.7 $\pm$ 12.3                             | 3.2 $\pm$ 0.8                                 |
| Biloruska Homostylina     | 2.2 $\pm$ 0.3                         | 24.9 $\pm$ 1.7                       | 86.5 $\pm$ 6.9                              | 2.9 $\pm$ 1.1                                 |
| Chornohlazka              | 2.0 $\pm$ 0.3                         | 25.6 $\pm$ 0.6                       | 77.0 $\pm$ 10.2                             | 3.6 $\pm$ 0.3                                 |
| Zhniarka                  | 2.1 $\pm$ 0.4                         | 25.7 $\pm$ 0.6                       | 80.2 $\pm$ 14.4                             | 4.1 $\pm$ 1.7                                 |
| Yampolska Mistseva        | 1.7 $\pm$ 0.4                         | 24.2 $\pm$ 0.9                       | 72.8 $\pm$ 20.8                             | 2.9 $\pm$ 1.1                                 |
| Yuvileina 100             | 1.5 $\pm$ 0.3                         | 27.4 $\pm$ 2.1                       | 54.1 $\pm$ 8.4                              | 3.2 $\pm$ 1.2                                 |
| Karmen                    | 1.6 $\pm$ 0.3                         | 26.8 $\pm$ 3.1                       | 58.6 $\pm$ 7.0                              | 3.0 $\pm$ 1.1                                 |
| Exida                     | 1.1 $\pm$ 0.4                         | 23.1 $\pm$ 0.5                       | 47.8 $\pm$ 19.0                             | 2.3 $\pm$ 1.4                                 |
| Anita Biloruska           | 1.8 $\pm$ 0.4                         | 25.5 $\pm$ 1.5                       | 69.2 $\pm$ 14.4                             | 2.5 $\pm$ 0.2                                 |
| Viktoriia Podilska        | 1.5 $\pm$ 0.7                         | 24.7 $\pm$ 1.4                       | 61.3 $\pm$ 24.8                             | 2.3 $\pm$ 0.8                                 |
| Roksolana                 | 1.4 $\pm$ 0.3                         | 30.7 $\pm$ 1.8                       | 46.2 $\pm$ 12.7                             | 2.2 $\pm$ 1.0                                 |
| Zhaleika                  | 1.3 $\pm$ 0.2                         | 24.9 $\pm$ 0.8                       | 50.5 $\pm$ 7.9                              | 2.7 $\pm$ 1.2                                 |
| Smuhlianka                | 1.7 $\pm$ 0.5                         | 26.7 $\pm$ 1.1                       | 64.9 $\pm$ 13.8                             | 3.6 $\pm$ 1.4                                 |
| Ketavase                  | 1.2 $\pm$ 0.4                         | 26.8 $\pm$ 2.2                       | 43.4 $\pm$ 9.7                              | 1.5 $\pm$ 0.6                                 |
| Kvietka Determinantna     | 1.4 $\pm$ 0.2                         | 28.3 $\pm$ 1.0                       | 49.8 $\pm$ 6.7                              | 2.8 $\pm$ 1.0                                 |
| Charovnitza               | 1.5 $\pm$ 0.7                         | 24.5 $\pm$ 1.8                       | 60.3 $\pm$ 29.8                             | 3.2 $\pm$ 1.7                                 |
| Sumchanka                 | 2.3 $\pm$ 1.1                         | 27.0 $\pm$ 0.7                       | 83.2 $\pm$ 39.7                             | 3.4 $\pm$ 1.6                                 |
| Bashkyrska Krasnostebelna | 1.4 $\pm$ 0.5                         | 24.5 $\pm$ 1.0                       | 57.7 $\pm$ 18.9                             | 3.9 $\pm$ 1.7                                 |
| Vlada                     | 1.8 $\pm$ 0.8                         | 27.7 $\pm$ 2.2                       | 61.0 $\pm$ 25.1                             | 4.6 $\pm$ 2.1                                 |
| Amethyst                  | 1.4 $\pm$ 0.6                         | 28.1 $\pm$ 1.7                       | 52.0 $\pm$ 22.1                             | 3.7 $\pm$ 1.8                                 |
| Lakneia                   | 1.5 $\pm$ 0.5                         | 28.1 $\pm$ 1.9                       | 54.0 $\pm$ 14.0                             | 3.9 $\pm$ 1.7                                 |
| Selianochka               | 1.9 $\pm$ 0.9                         | 29.6 $\pm$ 1.9                       | 65.7 $\pm$ 32.4                             | 3.4 $\pm$ 2.0                                 |
| Simka                     | 2.0 $\pm$ 0.6                         | 29.3 $\pm$ 1.1                       | 66.7 $\pm$ 19.2                             | 3.2 $\pm$ 1.5                                 |
| Oksana                    | 1.6 $\pm$ 0.8                         | 25.8 $\pm$ 2.4                       | 62.1 $\pm$ 29.8                             | 3.2 $\pm$ 1.9                                 |
| <b>Min</b>                | <b>1.1</b>                            | <b>23.1</b>                          | <b>43.4</b>                                 | <b>1.4</b>                                    |
| <b>Max</b>                | <b>3.2</b>                            | <b>31.5</b>                          | <b>112.7</b>                                | <b>4.7</b>                                    |
| <b>Mean</b>               | <b>1.7</b>                            | <b>27.0</b>                          | <b>63.5</b>                                 | <b>3.0</b>                                    |
| <b>Std. error</b>         | <b>0.1</b>                            | <b>0.4</b>                           | <b>2.8</b>                                  | <b>0.1</b>                                    |
| <b>Stand. dev</b>         | <b>0.5</b>                            | <b>2.0</b>                           | <b>15.3</b>                                 | <b>0.8</b>                                    |
| <b>Coeff. var</b>         | <b>26.2</b>                           | <b>7.5</b>                           | <b>24.1</b>                                 | <b>27.1</b>                                   |
| <b>Variance</b>           | <b>0.2</b>                            | <b>4.1</b>                           | <b>234.6</b>                                | <b>0.7</b>                                    |

\* $\pm$ SD – Standard Deviation



**Figure 4.** Dendrogram of cluster analysis (k-means clustering).

According to the obtained results, most of the studied buckwheat cultivars exhibit a high potential for productivity realization under the conditions of the Eastern Forest-Steppe of Ukraine, which is consistent with reports by other authors obtained in different climatic environments (Bisht et al., 2018; Kabanets et al., 2018; Amelin et al., 2020).

Clustering of the buckwheat cultivars was performed using the k-means method according to Ward's algorithm with Euclidean distance. Prior to the analysis, all traits were standardized using the z-score method. With the predefined number of clusters ( $k = 4$ ), the following clustering structure was identified.

Cluster 1 includes buckwheat cultivars with intermediate morphological and productivity traits. Plants in cluster 1 are characterized by medium plant height (91.2 cm), a moderate number of leaves (31.5 per plant), and inflorescences (22.9 per plant). Productivity per plant is the lowest among all groups (1.4 g), and thousand-kernel weight is also minimal (25.9 g). The number of kernels per plant and the seed set index are at intermediate levels. This cluster represents cultivars with a baseline level of agronomically valuable traits.

Cluster 2 includes cultivars exhibiting the shortest plant height (89.4 cm) and the lowest number of inflorescences (21.6 per plant). Despite these characteristics,



**Table 3.** Mean values of traits by cluster.

| Cluster   | Plant height, cm. | Number of leaves, pcs. | Number of inflorescences, pcs. | Productivity per plant, g | Thousand-kernel weight, g | Number of kernels per plant, pcs. | Number of kernels per inflorescence |
|-----------|-------------------|------------------------|--------------------------------|---------------------------|---------------------------|-----------------------------------|-------------------------------------|
| Cluster 1 | 91.2              | 31.5                   | 22.9                           | 1.4                       | 25.9                      | 53.9                              | 2.7                                 |
| Cluster 2 | 89.4              | 31.5                   | 21.6                           | 1.7                       | 27.0                      | 63.6                              | 3.3                                 |
| Cluster 3 | 91.9              | 36.7                   | 26.6                           | 2.4                       | 27.9                      | 85.7                              | 3.8                                 |
| Cluster 4 | 97.8              | 40.8                   | 30.5                           | 1.6                       | 27.4                      | 58.2                              | 2.1                                 |

productivity per plant is slightly higher than in Cluster 1 (1.7 g per plant), while the seed set index (3.3) and the number of kernels per plant (63.6) substantially exceed those of the first cluster. This may indicate a compact plant habit with relatively efficient yield formation.

Cluster 3 is the most valuable in terms of the analyzed traits. Cultivars in this group are characterized by the highest productivity per plant (2.4 g), the greatest number of kernels per plant (85.7), the highest thousand-kernel weight (27.9 g), and the best seed set index (3.8). At the same time, plant height and the number of vegetative organs remain at intermediate levels. This cluster represents an intensive type of cultivars and is the most promising for breeding purposes.

Cluster 4 includes cultivars with the most pronounced vegetative development. Plants in this cluster are tall (97.8 cm) and have the maximum number of leaves (40.8 per plant) and inflorescences (30.5 per plant). However, strong vegetative growth does not correlate with high productivity per plant (1.6 g), which remains low, while the seed set index (2.1) is the lowest among the identified clusters (Fig. 4, Table 3) (Bisht et al., 2018; Ghiselli et al., 2016).

## CONCLUSION

As a result of studying the buckwheat cultivar collection, significant genotypic variability was observed for

a complex of morphological and agronomically valuable traits under conditions in the Eastern Forest-Steppe of Ukraine. This confirms the expediency of using collection material of diverse ecological and geographical origins in breeding programs.

It was demonstrated that the formation of high buckwheat productivity is primarily determined by the number of kernels per plant and the level of seed set, both of which show a very strong positive correlation with productivity ( $r = 0.95$ ). Thousand-kernel weight is characterized by relative stability ( $V = 7.5\%$ ) and plays a stabilizing, rather than a determining, role in yield structure.

It was established that cultivars with a moderate number of inflorescences and better leaf surface supply per inflorescence (1.5–1.7 leaves per inflorescence) ensure more efficient realization of the generative potential of plants compared with tall forms exhibiting excessive vegetative development.

Based on the results of cluster analysis, a group of cultivars was identified that is characterized by maximum productivity per plant (2.4 g), the highest number of kernels per plant (85.7), high thousand-kernel weight (27.9 g), and a high seed set index (3.8). These cultivars are the most promising for breeding use. According to the study results, valuable sources of breeding traits include the cultivars 'Pidmoskovna T', 'M NF-84B', 'Mariia', 'Sumchanka', 'Roksolana', 'Krupynka', 'Selianochka', and 'Simka'.

## REFERENCES

- Alekseyeva, O. S., Taranenko, L. K., & Malina, M. M. 2004. Genetics, breeding, and seed production of buckwheat. Kyiv: Higher School, 213 p. [in Ukrainian]
- Amelin, A. V., Fesenko, A. N., Chekalin, E. I., Fesenko, I. N., & Zaikin, V. V. 2020. Higher yielding varieties of common buckwheat (*Fagopyrum esculentum* Moench) with determinate growth habit (single mutation det) manifest higher photosynthesis rate at stage of grain filling. Acta Agriculturae Slovenica, 115(1), 59–65. DOI: <http://dx.doi.org/10.14720/aas.2020.115.1.1316>

- Bhandari, H. R., Bhanu, A. N., Srivastava, K., Singh, M. N., & Shreya, H. A. 2017. Assessment of genetic diversity in crop plants-an overview. *Adv. Plants Agric. Res*, 7(3), 279–286. <https://doi.org/10.15406/apar.2017.07.00255>
- Bisht, A. S., Bhatt, A., & Singh, P. 2018. Studies on variability, correlation, and path coefficient analysis for seed yield in buckwheat (*Fagopyrum esculentum* Moench) germplasm. *Journal of Pharmacognosy and Phytochemistry*, 7(5S), 35–39.
- Cheadle, C., Vawter, M. P., Freed, W. J., & Becker, K. G. 2003. Analysis of microarray data using Z score transformation. *The Journal of molecular diagnostics*, 5(2), 73–81. [https://doi.org/10.1016/S1525-1578\(10\)60455-2](https://doi.org/10.1016/S1525-1578(10)60455-2)
- Chuiko, D., & Tryhub, O. 2025. Expression of Useful Traits in Determinant Buckwheat Accessions in the East of Ukraine. *Fagopyrum*, 42(1), 5–17. <https://doi.org/10.3986/fag0044>
- Dutta, P. P., Swain, S., Laloo, B., Kaur, S., Touthang, L., Verma, V. K., & Kumar, A. 2025. Ascertaining the morpho-molecular diversity in buckwheat germplasm and identification of high yielding, stable genotypes with superior biochemical quality. *Scientific Reports*, 15(1), 34232. <https://doi.org/10.1038/s41598-025-16156-5>
- Ermantaut, E. R., Goptsiy, T. I., Kalenska, S. M., Kryvoruchenko, R. V., Turchynova, N. P., & Prysyazhnyuk, O. I. 2014. Methodology of breeding experiments (in crop production). KhNAU named after V.V. Dokuchaev, 229 p. [in Ukrainian]
- Ghiselli, L., Tallarico, R., Mariotti, M., Romagnoli, S., Baglio, A. P., Donnarumma, P., & Benedettelli, S. 2016. Agronomic and nutritional characteristics of three buckwheat cultivars under organic farming in three environments of the Garfagnana mountain district. *Italian Journal of Agronomy*, 11(3), 729. <https://doi.org/10.4081/ija.2016.729>
- Hammer, Ø., & Harper, D. A. 2001. Past: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*, 4(1), 1–9.
- Hornýák, M., Słomka, A., Sychta, K., Dziurka, M., Kopeć, P., Pastuszak, J., ... & Płazek, A. 2020. Reducing flower competition for assimilates by half results in higher yield of *Fagopyrum esculentum*. *International Journal of Molecular Sciences*, 21(23), 8953. <https://doi.org/10.3390/ijms21238953>
- Jahn, R., Blume, H. P., Asio, V. B., Spaargaren, O., & Schad, P. 2006. Guidelines for soil description. FAO.
- Janovská, D., Jágr, M., Svoboda, P., Dvořáček, V., Meglič, V., & Hlásná Čepková, P. 2021. Breeding buckwheat for nutritional quality in the Czech Republic. *Plants*, 10(7), 1262. <https://doi.org/10.3390/plants10071262>
- Kabanets, V., Strakholis, I., & Klitsenko, A. 2018. Breeding of buckwheat varieties of different morphotypes and their distribution in Ukraine. *Bulletin of Agrarian Science*, 96(11), 141–146. [in Ukrainian]
- Kasajima, S., Namiki, N., & Morishita, T. 2016. Characteristics relating to the seed yield of determinate common buckwheat (*Fagopyrum esculentum* cv. Kitanomashu). *Fagopyrum*, 33, 1–5.
- Kharchenko, Yu. V., & Tryhub, O. V. 2018. Diversity of initial buckwheat material and directions of its use in breeding. *Genetic Resources of Plants*, (22), 31–43. DOI: 10.36814/pgp.2018.22.03 [in Ukrainian]
- Luthar, Z., Fabjan, P., & Mlinarič, K. 2021. Biotechnological methods for buckwheat breeding. *Plants*, 10(8), 1547. <https://doi.org/10.3390/plants10081547>
- Ohsawa, R. 2020. Current status and prospects of common buckwheat breeding in Japan. *Breeding science*, 70(1), 3–12. <https://doi.org/10.1270/jsbbs.19108>
- Pipan, B., Sinkovič, L., Neji, M., Janovská, D., Zhou, M., & Meglič, V. 2023. Agro-morphological and molecular characterization reveal deep insights in promising genetic diversity and marker-trait associations in *Fagopyrum esculentum* and *Fagopyrum tataricum*. *Plants*, 12(18), 3321. <https://doi.org/10.3390/plants12183321>
- Rauf, M., Yoon, H., Lee, S., Hyun, D. Y., Lee, M. C., Oh, S., & Choi, Y. M. 2020. Evaluation of *Fagopyrum esculentum* Moench germplasm based on agro-morphological traits and the rutin and quercetin content of seeds under spring cultivation. *Genetic Resources and Crop Evolution*, 67(6), 1385–1403. <https://doi.org/10.1007/s10722-020-00899-3>
- Shakalii, S. M., Yurchenko, S. O., Bahan, A. V., & Marinich, L. H. 2023. The influence of the variety on the manifestation of economic-valuable characters of buckwheat. *Scientific Progress & Innovations*, 26(2), 51–55. <https://doi.org/10.31210/spi2023.26.02.09>
- Shmaryayev, G. E., Yarchuk, T. Ya., & Yakushevsky, E. S. 1968. Methodological guidelines for studying maize, sorghum, and cereal crops (millet, buckwheat, rice) collection samples. Leningrad: VIR Printing House, 51 p. [in Russian]
- Suzuki, T., Noda, T., Morishita, T., Ishiguro, K., Otsuka, S., & Brunori, A. 2020. Present status and future perspectives of breeding for buckwheat quality. *Breeding Science*, 70(1), 48–66. <https://doi.org/10.1270/jsbbs.19018>

Volkodav, V. V. 2001. Methodology for examining varieties of cereals, pseudocereals, and legumes for suitability for distribution in Ukraine. Kyiv: UIESR, 112 p. [in Ukrainian]

## IZVLEČEK

### **Diferenciacija sort ajde (*Fagopyrum esculentum* Moench) glede na njihovo produktivnosti v gozdno-stepnem območju Ukrajine**

Predstavljeni so rezultati triletnih raziskav (2023–2025) 31 sort ajde (*Fagopyrum esculentum* Moench) različnih ekoloških in geografskih virov v Vzhodni gozdno-stepni Ukrajini. Avtorji so preučevali višino rastlin, število listov in socvetij, produktivnost rastlin, maso tisoč semen, število semen na rastlino in nastavitve semen. Ugotovljen je bil odločilen vpliv tipa rasti pri oblikovanju arhitekture rastlin in produktivnosti.

Ugotovljeno je bilo, da je produktivnost odvisna predvsem od števila semen na rastlino in nastavitve semen ( $r = 0.95$ ). Masa tisočih semen je relativno stabilna lastnost ( $V = 7.5\%$ ).

Sorte 'Pidmoskovna T', 'M NF-84B', 'Mariia', 'Sumchanka', 'Roksolana', 'Krupynka', 'Selianochka' in 'Simka' so bile prepoznane kot obetavni viri pomembnih lastnosti za nadaljnje križanje z namenom izboljšanja produktivnosti in teže tisoč zrn.